



Attitudes, Knowledge and Practice of Herbal Remedy Use among the Population Visiting Pharmacies and Health Care Providers in Alexandria, Egypt: A Cross-Sectional Study

Hoda Fathy^{1*}, Raghda El-Hawy² and Raghda Hassan²

¹Department of Pharmacognosy, Faculty of Pharmacy, Alexandria University, Egypt.

²Community Pharmacies, Alexandria, Egypt.

Authors' contributions

This work was carried out in collaboration among all authors. Author REH devised the idea for the study. Authors REH and RH analyzed the data. All authors were involved in collecting the data, interpreting the results of the analysis and critically reviewed the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2019/v30i230170

Editor(s):

- (1) Dr. Naseem A. Qureshi, National Center of Complementary and Alternative Medicine, Riyadh, Saudi Arabia.
(2) Prof. Marcello Iriti, Professor of Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

Reviewers:

- (1) Bagiu Radu-Vasile, Victor Babeş University of Medicine and Pharmacy, Romania.
(2) Danish Kadir, University of Rajshahi, Bangladesh.

Complete Peer review History: <https://sdiarticle4.com/review-history/52613>

Original Research Article

Received 09 September 2019
Accepted 11 November 2019
Published 15 November 2019

ABSTRACT

Aims: To show the levels of awareness, attitude, practices and socioeconomic factors related to consuming herbal products among Alexandrian citizen visiting pharmacies. The findings of this study are meant also to show the practices and beliefs of healthcare providers regarding using herbal remedies in treatment, including advising patients, reporting adverse effects and possessing knowledge about specific herb-drug interactions.

Study Design: people visiting pharmacies as well as pharmacists in their pharmacies and physicians in their clinics were invited to complete a questionnaire about their knowledge and personal experience in the consumption of herbal remedy. Anonymity was guaranteed. Questionnaire consists of three parts: demographic data, personal experience and a part concerning health care providers.

*Corresponding author: E-mail: hodasherif@alexu.edu.eg, hodasherif@hotmail.com;

Place and Duration of Study: Study area is the city of Alexandria, Egypt from the 1st July through September 2018.

Methodology: This is a descriptive cross-sectional study that used a self-administered, questionnaire from 213 participants (153+ 60 health care providers HCPs).

Results: Almost half the respondents preferred to be treated with herbs because they believed that herbs are safe. But they are not the proper choice to cure chronic diseases. Only 25% of herb users recorded suffering from side effects. The most common source of information about herbs was via internet, followed by family and friends, television and other types of media. Herbal remedies were purchased mainly from outlets other than pharmacies. The willingness to buy such products was not affected by participants' education level or their monthly income. The public didn't feel it's important to mention any herbal remedy they consume to their physician; similarly HCPs did not ask them. Generally, it was shown that HCPs have inadequate knowledge about herbs.

Conclusion: The prevalence of herb usage is moderate among the Alexandrian population. And it is imperative to educate HCPs about the benefits, risks and interactions of herbal supplements.

Keywords: Alexandria; herbal medicines; herb-drug interactions; questionnaire.

1. INTRODUCTION

During the last few decades, the phenomenon "back to nature" is spreading globally, along with a growing market demand for herbal remedies [1]. Even in industrialized regions such as Europe and North America, more than 50% of the population has used complementary or alternative medicine at least once [2]. According to the World Health Organization (WHO), about 80% of the population in most developing countries, especially in Africa, mainly depends on plant-based traditional medicines since they are often the only accessible and affordable treatment available for primary health care [3]. WHO also stated that health goals can't be achieved without the incorporation of herbal medicines [4].

Using herbs is common in the Arab world. Egypt has great plant biodiversity, which has provided a foundation for the use of traditional herbal medicine throughout the decades [4]. It is also important to note that medicinal herbs are not classified as drugs by the United State Food and Drug Administration (FDA) [5].

Until 2015, The Egyptian Guidelines for Registration of Herbal Medicines allowed for the free use of indigenous herbal remedies by the local community or in the local region, even if there was a lack of detailed information on these remedies. They had to meet the requirements of safety and efficacy, as specified in the Egyptian regulations for herbal medicines, only if they entered the market [6]. However, this changed in 2016, when traditional herbal medicines were defined as medicinal herbs used inside Egypt for a period of not less than fifteen years and that they should be supported by references for

safety and efficacy [7]. However, these regulations applied only to pharmacies, which are the only outlet under control of the Ministry of Health. In herb stores, herbal medicines are widely and freely available to Egyptian citizens, with no regulatory control [8].

Population in Egypt are accustomed to regularly consult pharmacist about minor as well as major health problems, they usually seek the pharmacy to buy their cosmetics "shampoo, diapers, creams and perfumes" the thing that made it easy to distribute the questionnaire in the pharmacy.

Health care providers have a professional responsibility to advise patients about potentially harmful aspects of herbal remedies, including contraindications or interactions with synthetic medications [9]. Previous studies from Nigeria, Kuwait, the United Arab Emirates and Australia have shown that most community pharmacists do not possess enough knowledge about the potential interactions and side effects of herbal medicines that they supply [10-12].

Thus, this study is designed to determine the awareness, patterns of use, attitude and socioeconomic factors among the general public visiting pharmacies and HCPs in Alexandria city, Egypt.

2. METHODOLOGY

2.1 General Procedures

2.1.1 Study area

Study area is the city of Alexandria, the second-largest city in Egypt situated on the northern

coast on the Mediterranean Sea, 2.6 Km², it has a large harbor and its population is more than four and half million (4,799,740 in March 2015) divided into 6 directorate .

2.1.2 Study design and setting

The study is based on a cross-sectional survey that used a self-administrated questionnaire designed to obtain information about the behavior regarding, knowledge of and use of medicinal herbs among Alexandrian society in different areas, with varied social and cultural levels.

We started with a pilot study on twenty Alexandrian citizens; not included in the study; to help determine the most suitable design and length of the questionnaire. Minor language adjustments were made.

2.1.3 Participants

Questionnaires were distributed and collected by the authors. Clients; eighteen years of age and above; who entered community pharmacies during the study period were enrolled in the study by way of a random sampling process—they were free to refuse participation in the survey. Data were collected anonymously via a self-administered questionnaire. Those who agreed to take part in the study were given the questionnaires, which were completed anonymously and collected after completion. They were assured of confidentiality, and they gave verbal consent to participate in the study.

2.1.3.1 Inclusion criteria

Clients; eighteen years of age and above; residents in Alexandria and are willing to participate in the questionnaire.

2.1.3.2 Exclusion criteria

- Non-Alexandrian citizens.
- Because it is self-administered questionnaire, Illiterate clients were not enrolled in the study.
- Clients under 18 years old.

2.1.4 Health care providers' participants

The questionnaire was handed to pharmacist in their pharmacies and to physicians in their

clinics. Both were free to refuse participation, and data collection was completed as above.

2.1.5 Questionnaire

The content of the questionnaire was generated by reviewing the literature [10,13–18]. No guidance or assistance was provided to participants as they completed the questionnaire during their visit to the pharmacy.

The questionnaire was in Arabic, and it comprised three sections. (the first two sections designed to be filled by both public respondents and HCP's and the third section directed only for HCP's) .The first section included questions related to sociodemographic and education characteristics, including age, gender, marital status, occupation, health insurance and monthly income. The second section addressed people's beliefs and behaviors related to herbal remedies. Respondents were asked to answer a series of questions concerning herbal remedy purchases and uses in the twelve months prior to the study, including the type of herbs being used, their main source ,their price, the reasons behind their use, and to report any adverse events they suffered. Participants were asked to provide only one answer for each question except for two questions; the first one regarding the herb source and the second whether they suffer from chronic disease. The third section had to do with HCPs' knowledge and how they advise their patients about the safe use of herbal products, how they report adverse effects, and how they check for an herb's interaction with conventional drugs that they dispense or prescribe.

The study was conducted from the 1st July through September 2018. An English translation of the questionnaire is available from the authors.

2.2 Statistical Analyses

The data were analyzed using the Statistical Package for the Social Sciences (SPSS) software program version 17.0 (SPSS Inc., Chicago, IL, USA).

3. RESULTS AND DISCUSSION

3.1 Demographic and Socioeconomic Data of the Study Population

In all, 153 participants, all residents of Alexandria, out of 200 distributed questionnaires in 6 different pharmacies each in one of the six directorate of Alexandria, responded and

completed the questionnaires appropriately for inclusion into the study. This was a response rate 76.5%.

In addition to a total of 60 out of 100 distributed questionnaires to HCPs (pharmacists and physicians) responded (60% response rate). The demographic and economic characteristics of the participants are listed in Table 1.

The majority of the healthcare respondents (66.3%) were between 25–39 years old, while 36.6% of the public respondents were between 40–60 years old. A total of 51 people reported having chronic disease.

The majority of total respondents (citizens and HCP's) were female (147, 69%). About two-thirds (63.8%) of the respondents had attended university. Reported monthly income varied, with 34.6% earning between 1,000–3,000 Egyptian pounds (LE), 15.1% earning more than 3000 LE, 9.7% earning between 500–1,000 LE, 20.5% with no income (housewives and students), and 14.1% preferring not to answer this question.

Most of the respondents were married (115 participants, 56%), and 65.2% (139 participants) had health insurance. 63.7% of citizens participants were government employees.

3.2 The Use of Herbal remedies

A total of 109 (93 + 16 HCPs) preferred to use herbs in general for treatment. But only 98 (76 + 22 HCP's) of them admitted to prior treatment with herbs—35 were males, and 63 were females-.

3.3 The Reason behind choosing to be Treated with Herbal Remedies

Overall, 89 (77 + 12) HPCs, from the 109 herb users, believed that herbs are safe. The details of the remaining reasons are summarized in Table 2.

3.4 Source of Herbs

The 98 that were previously treated with herbs admitted that the most common source for buying herbs was at herb stores (60 [50 +10 HCPs]), followed by 24 (17 + 7 HCPs) from pharmacies, 6 from supermarkets, 6 from family or friends and 2 from TV commercials. Strikingly, when linking the educational levels of herb users

to the source of buying herbs, 33 of the university graduates bought the herb from herb stores, versus only 6 buying from the pharmacy. The monthly income did not affect the source of buying the herbs.

3.5 Side Effects Reported

Few (26%) herb users indicate that they have experienced certain associated side effects. Six participants suffered from nausea, two from vomiting, seven from diarrhea, four from constipation, one from irritability, one from skin rash and four didn't specify which side effects. It is worth mentioning that 15 out of 21 stated that they obtain their information about herbs from the internet, and eight said that they get their information from family and friends. Unfortunately, no one mentioned if they reported these side effects to a physician. In addition, no one specified the name of the herb(s) that caused these side effects.

3.6 The Reasons that Encourage them to Buy A Herb

These are summarized in Table 3.

3.7 The Maximum Price they are willing to pay for a Herb

The study showed that the maximum limit that participants will pay to purchase herbal products ranges between 1–10 LE equivalent to 0.06-0.6 USD (29.5% [53 + 10 HCPs]) and 11–50 LE equivalent to 0.68- 3.1 USD (35.6% [54 + 25 HCPs]). A total of 28 participants were willing to pay between 51–100 LE equivalent to 3.18-6.2 USD, and the remaining 10 chose to spend between 100–300 LE equivalent to 6.2-18.7 USD. It is worth mentioning that 14 of those 28 stated that their monthly income is above 3,000 LE (187 USD).

3.8 Effect of Monthly Income on Using Herbs

Monthly income did not affect the decision on whether to use herbs, as the ratio between herb users and non-users was almost the same among all categories.

3.9 Source of Information about the Herbs

The most common sources for recommending and providing information regarding the use of herbs was the internet followed by family and friends, Table 4.

Table 1. Demographic and socioeconomic data of study population

	Frequency of citizens (153)	Frequency of HCP's (60)	Total
Sex			
Male	57	8	65
Female	95	52	147
missing	1	-	1
Age			
18-24	42	13	55
25-39	46	40	86
40-60	56	6	62
Above 60	8		8
missing	1	1	2
Marital status			
Single	56	32	88
Married	52	16	68
Married with children	38	9	47
Divorced or widow	8	3	11
Working status			
None	8		
Government employee	88		
Worker	2		
housewife	15		
student	35	7 (postgraduate)	
other	5	53	
Educational level			
Non educated	1		1
Primary	4		4
Middle school	30		30
High school	16		16
University	95	41	136
Postgraduate	7	13	20
Monthly income 'L.E.			
none	42		42
500-1000 (30-60 'USD)	19	1	20
1000-3000 (60-180 USD)	40	31	71
Above 3000 (above 180 USD)	26	4	31
Prefer not to answer	26	24	51
Chronic diseases*			
None	-	48	
Hepatic diseases	-	-	
Renal diseases	107		115
Diabetic	2		2
Cardiac diseases and hypertension	10		10
Bronchial asthma	10		10
Bronchial asthma	24	1	24
Others	3	2	3
missing		9	
Health insurance			
Yes	100	39	139
No	52	13	65
missing	1	7	8

*participants were allowed to choose more than one disease
'L.E is the acronym of Egyptian pound, USD united state dollars

Table 2. Frequency of herb users and non users and reason behind their opinion

Reason	Herb users		Reason	Herb non users	
	Frequency	HCP's		Frequency	HCP's
Herbs are safe	77	12	I didn't get the expected results	14	14
They have rapid onset of action	4	1	I don't have a trusted source of information about herbs	34	23
Their reasonable price	5	2	Herbs might have dangerous side effects	5	5
Herbs are more effective than regular medications	7	-	Their price is not reasonable	1	-
Missing	-	1		3	2
Total	93	16		60	44

Table 3. The frequency of the reasons that encourage them to buy a herb

Reason*	Frequency	Frequency of HCPs
Price	50	9
As a nice drink	72	27
According to speed of onset of action	19	8
Package	2	1
I trust the store	16	8
Whether there is any associated side effects	15	12
Media	30	12
Known brand	27	14
Good offer	3	-

*participants were allowed to choose more than one reason

Table 4. Participants' opinions regarding the current available source of information on herbs

Source	Frequency
Television and media	47
Internet	89
Family and friends	55
Herb stores	25
Pharmacist	33
Physician	12
others	5

The demographic and economic characteristics of the participants are listed in Table 1. The majority of the healthcare respondents (66.3%) were between 25–39 years old, most of them were females (52, 86.7%) and single (32 participants, 53.3%). with only 3 reported having chronic disease. Their monthly income varied, with 31 earning between 1,000–3,000 Egyptian pounds (LE), 4 earning more than 3000 LE, 1 earning between 500–1,000 LE, and 24 preferring not to answer this question.

3.10 Whether Herb Can Treat Chronic Diseases

A total of 75 (62 + 13 HCPs) believed that they can, while 134 (90 + 44 HCPs) believed that they cannot. An education level up to university graduates and postgraduates (almost 60 [70%]) were more likely to believe that they cannot.

16 preferred to use herbs in general for treatment. But 24 admitted to prior treatment with herbs. The most common source for obtaining herbs was at herb stores 10, followed by 6 from pharmacies, 4 from a physician, and 6 from supermarkets. Six herb users suffered from nausea, two from vomiting, seven from diarrhea, four from constipation, one from irritability, one from skin rash and four from unspecified side effects.

3.11 Health Care Providers' Responses

The following is the summary of their answers on the first and second sections:

Among the non users, 23 were concerned about the source of the herbs, 5 were afraid they might suffer from dangerous side effects, 14 stated that they did not see any results when using herbs,

and the remaining were worried about herb prices.

3.12 The Third Sector of the Questionnaire

This sector was designed to target the health care providers; pharmacists and physicians; to measure the extent of their behavior and knowledge relating to the study topic.

3.12.1 Attitude towards herbal remedy

The first question was “Before prescribing a medication, do you ask the patients if they are taking medicinal herbs?” The answers differed between “I occasionally ask them” (23 [43.4%]), “No, I do not ask them” (23 [43.4%]) and “I often ask them” (7 [13.2%]).

3.12.2 Dispensing/prescribing herbal remedies

A total of 71% did not prefer to dispense/prescribe herbs for treatment, and 78% did not believe that herbs can treat chronic diseases. The reasons behind their refusal to use herbs included not getting the required results (26%) or concerns about the herb source (38.5%).

The third question, “Do you advise your patients to consume herbs as a medication?” was crucial to figure out HCPs’ behavior regarding the dispensing of medicinal herbs. The participants answered evenly between “yes” (50%) and “no” (50%).

3.12.3 Source of information about herbs

Almost 30% chose the most commonly trusted sources to provide information, including “by Google search” (28%), “searching in books” (30%), and asking academic colleagues or a pharmacist (18%). The worrisome percentages were, 12% get their information from family and friends, 6% get their information from herb stores, and a small percent do not bother to ask (6%).

3.12.4 General knowledge about herb-drug interactions

At the end of the questionnaire, we added three questions that may give us a general perspective about medical participants’ knowledge of medicinal herbs. The first question was: “What is the herbal product that interacts with

hypolipidemic drugs such as Atorvastatin”. They were offered three choices; garlic, ginkgo and hibiscus. The correct answer was “garlic,” which has a powerful synergistic effect with hypolipidemic drugs [18]. A total of 57.7% of respondents answered this question correctly, and 11 participants didn’t respond to the question.

The second question was about the herb that may cause abortion or premature labour to pregnant women. They had three answers to choose from fenugreek, parsley or anise and the correct answers were “fenugreek” or “parsley”—potent uterine stimulants [19,20]. The percentage of those who answered correctly was 59.6%. Five chose not to answer, and three chose “anise”.

The third and final question was about the herb-drug interaction between the evening primrose herb with anticoagulants, blood pressure medications and non steroidal anti-inflammatory drugs [21]. A total of 25% answered correctly, with eight participants didn’t answer.

4. CONCLUSION

Most of the studies conducted in the United States and Europe were interested in complementary medicine in general, a term that encompasses many forms of treatment, such as acupuncture and spiritual healing. Only few of the studies were focused purely on the use of herbal remedies. Rates of use were low, ranging between 10–20%, whereas in the United Arab Emirates, the percentage of herbal remedy users was higher (60%), compared to 43% in this study. This was partly expected, due to the different cultures of the Arab world and Western countries [22]. Herb users believed that herbs are safe, which agrees with worldwide popular belief [14].

Meanwhile, the need for accurate, up-to-date information on the prevalence and socio-cultural and personal factors (knowledge, beliefs and attitudes) that underlie an individual’s decision to use herbal remedies is of national importance.

Study evidence shows that many herbal remedies are mainly purchased from outlets other than pharmacies, where herbal remedies are sold freely, without regulation of the Ministry of Health [21]. Herbal stores in Egypt are usually run by unqualified personnel, who transmit the information from generation to generation. Herbs

are vulnerable to being contaminated, misidentified or adulterated.

Most of the information about herbs is taken from television and the internet due to the increased online availability of information about herbal remedies [22]. Both the public and their HCPs do not give/ask information about any herbal remedy they use when prescribing or during consuming conventional drugs. Nondisclosure of herbal remedy use may cause individual to be at risk of undue harm. This may be justified by Fear of negative response from doctors. In addition to doctor's communication skills and providing enough time for the patient [23]. A study conducted in USA showed another perspective which is; most of the physicians were not comfortable in counseling their patients about CAM treatments [24].

Although natural health products are routinely available without prescription, that does not mean they are completely safe for all individuals. Many of the participants (99, 46.5%) admitted consuming common herbs daily, as soothing drinks, without knowing that these are actually herbal drugs and without knowledge of possible serious drug interactions [25].

Few (26%) herb users indicate that they have experienced certain associated side effects, which is higher than the number reported in North America [17]. Monthly income did not affect the decision on whether to use herbs, as the ratio between herb users and non-users was almost the same among all categories.

Social influencing factors, such as friends or family members, have been reported in other studies and are consistent with the results of the factor analysis of this questionnaire [26].

The findings related to the influence of sex on herbal remedy use were found to be slightly higher percentage of women compared to men in agreement with the previous study [26,27]. The reasons for consuming herbs ranged from safety, perceived efficacy and ease of access—this agrees with previous studies [26]. In addition, the study showed a deficit in information among HCPs regarding herb-drug interactions. Knowledge deficiencies found in this study were also reported by many studies in the Arab region, including Saudi Arabia, Jordan, Oman, Kuwait, Qatar and Lebanon, as well as in the United States [26, 28,29]. A possible reason for HCP's missing answers is because participants were

hesitant to answer them, as they did not know the correct answers [28-30].

There is a need of proper education about herbal products among pharmacists, especially community pharmacists, who are expected to provide correct information about their proper use, adverse effects and interactions [29-30].

5. POTENTIAL LIMITATIONS TO THIS STUDY

These include the small number of responders; the majority of whom were females may be because women are more open and receptive to a health concern and tend to share problems [23,27]. Future studies should focus on testing a larger sample, of equal males and females percentages in order to avoid gender bias. The self-administered questionnaire also has limitations, as the data collection relied on self-reported answers, which could be subject to errors because of memory recall or social-desirability bias.

CONSENT

All authors declare that 'verbal informed consent was obtained from the participants.

ETHICAL APPROVAL

The survey was approved by the ethical committee in the faculty of pharmacy, Alexandria university, Egypt and was given the number (ec18/1) confirms either that this study is not against the public interest, or that the release of information is allowed by legislation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2014;4:177. DOI: 10.3389/fphar.2013.00177
2. Al-Kindi RM, Al-Mushrafi M, Al-Rabaani M, Al-Zakwani I. Complementary and alternative medicine use among adults with diabetes in Muscat Region, Oman. *Sultan Qaboos Univ Med J.* 2011;11(1):62–68. DOI: org/10.15344/2456-8171/2016/106

3. WHO traditional medicine strategy 2014-2023.
Available:https://www.who.int/medicines/publications/traditional/trm_strategy14_23/en/
4. Mostafa N, Singab A. Prospective of Herbal Medicine in Egypt. *Med Chem (Los Angeles)*. 2018;8:116-117.
DOI: 10.4172/2161-0444.1000502
5. Dietary supplement Health and Education Act 1994, Public Law 103-417,103rd congress page. Food and Drug Administration Website.
Available:<http://www.fda.gov/opacom/laws/dshea.html>. [Access date 05.03.2019]
6. The Egyptian Guidelines for Registration of Herbal Medicines; 2015.
Available:http://www.eda.mohp.gov.eg/images/News/F_53.pdf
[access date 05.06.2019]
7. The Egyptian Guidelines for Registration of Herbal Medicines; 2016.
Available:http://www.eda.mohp.gov.eg/images/News/F_67.pdf
[Access date 05.06.2019]
8. Pharmaceutical country profile.
Available:https://www.who.int/medicines/areas/coordination/Egypt_PSCPNarrativeQuestionnaire_27112011.pdf.
[access date 29.05.2019]
9. Farrell J, Ries N, Boon H. Pharmacists and Natural Health Products: A systematic analysis of professional responsibilities in Canada. *Pharm Pract (Granada)*. 2008;6(1):33–42.
DOI: 10.4321/s1886-36552008000100006
10. Oshikoya K, Oreagba I, Ogunleye OO, Oluwa R, Senbanjo I, Olayemi S. Herbal medicines supplied by community pharmacies in Lagos, Nigeria: pharmacists' knowledge. *Pharm Pract (Granada)*. 2013;11(4):219–227.
DOI: 10.4321/s1886-36552013000400007
11. Hutchinson S, Mitchell K, Hansford D, Stewart D. Community pharmacists' views and experiences of counter-prescribing in pregnancy. *Int J Pharm Pract*. 2001;9:15–21;
DOI:org/10.1111/j.2042-7174.2001.tb01024.x
12. Chang ZG, Kennedy DT, Holdford DA, Small RE. Pharmacists' knowledge and attitudes toward herbal medicine. *Ann Pharmacother*. 2000;34:710–715.
DOI: 10.1345/aph.19263
13. Sekhria K, Bhanwrab S, Nandhaa R. Herbal products: A survey of students' perception and knowledge about their medicinal use. *Int J Basic Clin Pharmacol*. 2013;2:71-6.
DOI: 10.5455/2319-2003.ijbcp20130114
14. Griffin B., Citkovitz C. Survey: Preferences and Limitations of Herbal Medicine Use Among Patients at a Community Acupuncture Clinic. *Acupunct Med*. 2017; 29(1).
DOI: org/10.1089/acu.2016.1204
15. Ameade E., Amalba A., Helegbe G. Mohamed B. Herbal medicine: a survey on the knowledge and attitude of medical students in Tamale, Ghana model. *PJMPr*. 2015;3(1)1-8.
DOI: 10.1016/j.jtcme.2015.03.004
16. Fahmy SA, Abdu S, Abuelkhair M. Pharmacists' attitude, perceptions and knowledge towards the use of herbal products in Abu Dhabi, United Arab Emirates. *Pharm Pract (Granada)*. 2010; 8(2):109-115.
DOI: 10.4321/S1886-36552010000200005
17. Awad A. and Al-Shaye D. Public awareness, patterns of use and attitudes toward natural health products in Kuwait: a cross-sectional survey. *BMC Complem Altern M*. 2014;14:105.
DOI: 10.1186/1472-6882-14-105
18. Reddy GD, Reddy AG, Rao GS, Kumar MV. Pharmacokinetics interaction of Garlic and atorvastatin in dyslipidemic rats. *Indian J. Pharmacol*. 2012;44(2):246-252.
DOI: 10.4103/0253-7613.93860
19. Sim T, Sheriff J, Hattingh L, Tee L. The use of herbal medicines during breastfeeding: a population-based survey in Western Australia. *BMC Complem Altern*. 2013;13:317.
DOI.org/10.1186/1472-6882-13-317
20. Shinde P, Patil P. and Bairagi V. Herbs in pregnancy and lactation: a review appraisal. *IJPSR*, 2012; Vol. 3(9): 3001-3006.
DOI.org/10.13040/IJPSR.0975-8232
21. Available:WebMd.www.content.health.msn.com/content/drugs/2/4046
[Accessed March 10, 2019]
22. AlBraik A, Rutter M, Brown D. A cross-sectional survey of herbal remedy taking by United Arab Emirate (UAE) citizens in Abu Dhabi. *Pharmacoepidemiol drug saf*. 2008;(7): 725-32.
DOI: 10.1002/pds.1591
23. Kelak JA, Cheah WL, Safii R. Patient's decision to disclose the use of traditional and complementary medicine to medical doctor: A descriptive phenomenology

- study. Evid Based Complement Alternat Med. 2018;4735234.
DOI: [org/10.1155/2018/4735234](https://doi.org/10.1155/2018/4735234)
24. Wahner-Roedler DL, Vincent A, Elkin PL, Loehrer LL, Cha SS, Bauer BA. Physicians' attitudes toward complementary and alternative medicine and their knowledge of specific therapies: a survey at an academic medical center. Evid Based Complement Alternat Med. 2006;3(4):495–501.
DOI: [10.1093/ecam/nel036](https://doi.org/10.1093/ecam/nel036)
25. Sharma V, Holmes JH, Sarkar IN. Identifying Complementary and Alternative Medicine Usage Information from Internet Resources. A Systematic Review. Methods Inf Med. 2016;55(4):322–332.
DOI: [10.3414/ME15-01-0154](https://doi.org/10.3414/ME15-01-0154)
26. Patterson C, Arthur H. A complementary alternative medicine questionnaire for young adults. Integr Med Insights. 2009;4:1–11.
DOI: [.org/10.4137/IMI.S2281](https://doi.org/10.4137/IMI.S2281).
27. Alwhaibi M, Sambamoorthi U. Sex Differences in the Use of Complementary and Alternative Medicine among Adults with Multiple Chronic Conditions. Evid Based Complement Alternat Med. 2016;2067095.
DOI: [10.1155/2016/2067095](https://doi.org/10.1155/2016/2067095)
28. Hijazi MA, Shatila H, El-Lakany A, et al. Beliefs, Practices and knowledge of community pharmacists regarding complementary and alternative medicine: national cross-sectional study in Lebanon. BMJ Open. 2019;9:e025074.
DOI: [10.1136/ bmjopen-2018-025074](https://doi.org/10.1136/bmjopen-2018-025074)
29. Hilal M, Hilal S. Knowledge, attitude and utilization of herbal medicines by physicians in the Kingdom of Bahrain: A cross-sectional study. J Assoc Arab Univ Basic App Sci. 2017;24:325-333.
DOI: [.org/10.1016/j.jaubas.2016.11.001](https://doi.org/10.1016/j.jaubas.2016.11.001).
30. Harris IM, Kingston RL, Rodriguez R, et al. Attitudes towards complementary and alternative medicine among pharmacy faculty and students. Am J Pharm Educ. 2006;70:129.
DOI: [10.5688/aj7006129](https://doi.org/10.5688/aj7006129)

© 2019 Fathy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://sdiarticle4.com/review-history/52613>